



TUMORIGENESIS AND NEOPLASTIC PROGRESSION

Fibrosis and Hypoxia-Inducible Factor-1 α — Dependent Tumors of the Soft Tissue on Loss of Von Hippel-Lindau in Mesenchymal Progenitors



Laura Mangiavini,^{*†‡§} Christophe Merceron,^{*†¶||} Elisa Araldi,[‡] Richa Khatri,[‡] Rita Gerard-O'Riley,[†] Tremika L. Wilson,^{*†‡} George Sandusky,^{**} Jerome Abadie,^{††} Karen M. Lyons,^{‡‡} Amato J. Giaccia,^{§§} and Ernestina Schipani^{*†‡}

From the Department of Orthopaedic Surgery and the Division of Endocrinology, Department of Medicine,* University of Michigan Medical School, Ann Arbor, Michigan; the Division of Endocrinology,[†] Department of Medicine, and the Department of Pathology,** School of Medicine, Indiana University, Indianapolis, Indiana; the Endocrine Unit,[‡] Massachusetts General Hospital-Harvard Medical School, Boston, Massachusetts; the Department of Orthopaedic and Traumatology,[§] Milano-Bicocca University, Milan, Italy; INSERM,[¶] UMRS 791-LIOAD, Centre for Osteoarticular and Dental Tissue Engineering, Group STEP (Skeletal Tissue Engineering and Physiopathology), Nantes, France; the Faculty of Dental Surgery,^{||} and Oniris Animal Cancers, Models for Comparative Oncology Research (AMaROC),^{††} l'Universite Nantes Angers le Mans (LUNAM), Nantes, France; the Department of Orthopaedic Surgery,^{‡‡} David Geffen School of Medicine at the University of California Los Angeles, Los Angeles, California; and the Division of Radiation and Cancer Biology,^{§§} Department of Radiation Oncology, Center for Clinical Sciences Research, Department of Radiation Oncology, Stanford University, Stanford, California

Accepted for publication
July 27, 2015.

Address correspondence to Ernestina Schipani, M.D., Ph.D., University of Michigan, Medical School, A. Alfred Taubman Biomedical Science Research Bldg., 109 Zina Pitcher Place, Room 2007, Ann Arbor, MI 48109; or Amato J. Giaccia, Ph.D., Division of Radiation and Cancer Biology, Department of Radiation Oncology, Stanford University, CCSR-South, Room 1255, 269 Campus Dr., Stanford, CA 94303-5152; or Karen M. Lyons, Ph.D., Department of Orthopaedic Surgery, David Geffen School of Medicine, University of California, Los Angeles, CA 90095. E-mail: eschipani@med.umich.edu, giaccia@stanford.edu, or klyons@mednet.ucla.edu.

The hypoxia-inducible factor (Hif)-1 α (Hif-1 α) and Hif-2 α (Epas1) have a critical role in both normal development and cancer. von Hippel Lindau (Vhl) protein, encoded by a tumor suppressor gene, is an E3 ubiquitin ligase that targets Hif-1 α and Epas1 to the proteasome for degradation. To better understand the role of Vhl in the biology of mesenchymal cells, we analyzed mutant mice lacking Vhl in mesenchymal progenitors that give rise to the soft tissues that form and surround synovial joints. Loss of Vhl in mesenchymal progenitors of the limb bud caused severe fibrosis of the synovial joints and formation of aggressive masses with histologic features of mesenchymal tumors. Hif-1 α and its downstream target connective tissue growth factor were necessary for the development of these tumors, which conversely still developed in the absence of Epas1, but at lower frequency. Human tumors of the soft tissue are a very complex and heterogeneous group of neoplasias. Our novel findings in genetically altered mice suggest that activation of the HIF signaling pathway could be an important pathogenetic event in the development and progression of at least a subset of these tumors. (*Am J Pathol* 2015, 185: 3090–3101; <http://dx.doi.org/10.1016/j.ajpath.2015.07.008>)

The ability of a cell to adapt to low oxygen tension (hypoxia) is critical in normal development and differentiation and in pathologic settings, such as cancer and ischemia. The hypoxia-inducible factor (Hif)-1 α and Hif-2 α (Epas1) are crucial mediators of the homeostatic responses that allow hypoxic cells to survive and differentiate.¹ In normoxia, the E3 ubiquitin ligase von Hippel Lindau (Vhl) binds to Hif-1 α and Epas1 and targets them to the proteasome for degradation.^{2–4} Cells that lack functional Vhl are unable to degrade the Hif- α s, and this ultimately results in their progressive accumulation.^{2,3,5,6} However, Vhl, which is a ubiquitously expressed protein, has also a variety of biological activities,

including control of the cell cycle, regulation of matrix proteins, and interaction with the cytoskeleton and the primary cilia that are HIF independent.^{7,8}

Notably, VHL is a tumor suppressor, and its deficiency is a key pathogenetic event in a variety of human familial and

Supported by NIH grant RO1 AR065403 (E.S. and A.J.G.) and grant FP7/2007-2013 from the People Program (Marie Curie Actions) of the European Union's Seventh Framework Program registered under the Research Executive Agency grant agreement 300388 (C.M.).

L.M. and C.M. contributed equally to this work.

Disclosures: None declared.

sporadic tumors.⁹ However, to this end, it is still largely unknown whether HIF-dependent or HIF-independent VHL functions are critical for VHL-defective tumorigenesis.⁸ Moreover, the phenotypic identity of the cells that give origin to VHL-deficient tumors is also the object of intense debate. Along these lines, in VHL-deficient hemangioblastomas, genetic alterations consistent with a neoplastic nature have been identified in the mesenchymal/stromal component of these tumors.^{10,11} Moreover, a mesenchymal cell origin has been provocatively suggested for VHL-associated clear cell renal carcinomas.¹²

We previously reported that Vhl is critically important for the proper development of cartilage and bone.^{13–16} To better understand the role of Vhl in the biology of mesenchymal cells, we analyzed mutant mice that lacked Vhl in mesenchymal progenitors that give rise to the soft tissues that form and surround synovial joints. We thus found that Vhl prevents the development of fibrosis and mesenchymal tumors at these sites by regulating HIF activity.

Materials and Methods

All the experiments were performed using at least three independent biological replicates.

Generation of Mice

Generation and genotyping of the *Vhl* (FVB/N), *Hif1a* (FVB/N), *Epas1* (C57/B6), floxed mice, *Prx1-Cre* (FVB/N), *Col2a1-Cre* transgenic mice, and *ROSA26 mT/mG* (FVB/N) reporter mice have been previously described.^{15,17–22} Connective tissue growth factor (*Ctgf*) (C57/B6) floxed mice were generated and characterized in the laboratory of Dr. K.M. Lyons.²³ Further details on *Ctgf* floxed mice are given elsewhere. For all studies, *cre*-positive heterozygous floxed mice and *cre*-negative homozygous floxed mice, respectively, were used as controls.

Cartilage and bone phenotype of VHL, VHL–HIF-1, and VHL–HIF-2 mice have been previously described.¹⁶

All procedures that involved mice were performed in accordance with the NIH guidelines for use and care of live animals and were approved by University of Michigan Committee on Use and Care of Animals (protocol PRO00005182).

Cryosections, Routine Histologic Analysis, Terminal Deoxynucleotidyl Transferase-Mediated dUTP Nick-End Labeling, *In Situ* Hybridization, and Immunohistochemistry Detection on Paraffin Sections

Frozen sections from p14 hindlimbs and sternum were prepared and cut as previously described.¹⁶ Frozen sections from p21 COL2Cre-mTmG hindlimbs were prepared according to Kowamoto Tape Transfer Kit (SECTION-LAB Co. Ltd., Hiroshima, Japan).^{24,25} Briefly, freshly isolated hindlimbs were embedded in Super Cryo Embedding Medium and




Figure 1 *Cre* is active in the soft tissues surrounding bone and cartilage of PRX1-Cre mice. **A–D**: Detection of fluorescence in frozen sections of tibias isolated from postnatal day 14 PRX1Cre-mTmG mice (**A**). Closeups of the synovium (**middle boxed area**; **B**), perimysium and epimysium (**top boxed area**; **C**), and one blood vessel (**bottom boxed area**; **D**) are shown. The enhanced green fluorescent protein signal indicates tissues and cells in which *cre* recombination occurred. Scale bars: 500 μ m (**A**); 100 μ m (**B–D**).

frozen in cold hexane. Frozen blocks were positioned in the cryostat, and a double-sided tape provided by the kit was applied on the block. Ten-micrometer sections were then cut with tungsten carbide blades and stored at -80°C .

Routine histologic analysis, terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling assay, and *in situ* hybridization on embryonic stages E15.5 and E17.5, newborn and postnatal ages p13–p28 forelimbs and hindlimbs were performed as previously reported.^{16,26}

For immunohistochemistry (IHC), paraffin sections from forelimbs of p21 to p28 mice were treated with sodium citrate buffer (pH 6) at 95°C for 15 minutes. Sections were then incubated with the following primary antibodies: anti-cadherin-11 at 1:100 (71-7600, Invitrogen, Life Technologies, Grand Island, NY), anti-Ctgf at 1:100 (sc-14939, Santa Cruz Biotechnology, Inc., Dallas, TX), anti-Vhl (BD556347, BD Biosciences, San Jose, CA) overnight at 4°C , anti-CD31 at a 1:25 (DIA310, Dianova, Hamburg, Germany), anti-desmin at 1:200 (AF3844, R&D Systems, Minneapolis, MN), anti-S100 at 1:2500 (Z0311, Dako North America, Inc., Carpinteria, CA), anti-cytokeratin at 1:100 (M0821, Dako North America), anti-vimentin at 1:200 (M7020, Dako North America), anti-von Willebrand factor at 1:200 (A0082, Dako North America) for 30 minutes at room temperature, and anti-CD45 at 1:200 (BD550539, BD Biosciences) for 1 hour at room temperature. After incubation with the




Figure 2 Deficiency of von Hippel Lindau (Vhl) in limb bud mesenchyme causes fibrosis of the synovial joints. Hematoxylin and eosin staining of postnatal day (p) 20 shoulders (A and B), p17 elbows (C and D), p17 wrists (E and F), and p18 knees (G and H) in control (CNTRL) (A, C, E, and G) and VHL (B, D, F, and H) mice. The **arrows** indicate the fibrosis (B) and point to the fibrotic infiltration (D). **F**: The **white arrow** indicates the fusion of small cartilaginous elements and the **black arrow** indicates the ectopic cartilage. The mutant knee joint is occupied by a massive fibrotic infiltration in **H**. Scale bar = 100 μ m.

appropriate biotinylated secondary antibody, binding was detected using the labeled streptavidin biotin and the tyramide amplification system (Perking Elmer, Waltham, MA) following the manufacturer's instructions.

Image Acquisition

Images were acquired with Eclipse E800 (Nikon, Tokyo, Japan). Additional images were captured with a Leica DM LB compound microscope (Leica Microsystems, Wetzlar, Germany). For fluorescent images, frozen sections were dehydrated at 4°C overnight, stained with DAPI (Life Technologies), and then overlaid with coverslips onto ProLong Diamond

Antifade Mountant (Life Technologies). Selected photographs were taken using a Nikon A1 confocal microscope (Nikon) with filters for red fluorescent protein, fluorescein isothiocyanate, and DAPI. Representative images of each technique are shown.




Figure 3 Aggressive synovial fibrosis in shoulder and elbow of von Hippel Lindau (VHL) mutants. **A–C**: Hematoxylin and eosin (H&E) staining of postnatal day (p) 18 scapulas in control (CNTRL) (A) and VHL (B) mice. **C**: Higher magnification of **B**. Aggressive fibrotic lesions are present in the shoulder of mutant mice. **D**: *In situ* hybridization for detection of Col1a1 mRNA of p18 scapula in VHL mice. These fibrotic lesions express Col1a1 mRNA. **E** and **F**: Radiographs of p21 forelimbs in CNTRL (E) and VHL (F) mice. The **white arrows** (F) indicate the erosions of the ulna. The **black arrow** (F) points to the persistent deformity of the humerus observed in VHL mice. **G** and **H**: H&E staining of p20 elbows in CNTRL (G) and VHL (H) mice. The **black arrows** (H) highlight the infiltration of the aggressive fibrosis. Scale bars: 100 μ m (A–D, G, and H); 0.5 cm (E and F).




Figure 4 Presence of tumors in forelimbs and hindlimbs of von Hippel Lindau (VHL) mice. The aggressive fibrosis postnatally was associated with the formation of tumors. The development of these tumors mainly occurred in the scapula and elbow, and, sporadically, also in the knee. Radiographs revealed that the VHL masses were partially mineralized and were highly heterogeneous with irregular borders. Radiographs of postnatal day (p) 21 to 28 elbows (A–E), p21 to 28 shoulders (F–J), and p21 knees (K and L) in control (CNTRL) (A, F, and K) and VHL (B–E, G–J, and L) mice. Scale bar = 0.5 cm.

Radiography

Forelimbs were analyzed by radiography, using an XPERT 80- L Cabinet X-ray System (Kevex-90Kv; Kubtec X-ray, Milford, CT) as previously reported.¹⁶ In addition, some forelimbs were analyzed using a Faxitron X-ray cabinet (Faxitron Bioptics, Tucson, AZ) as previously reported.²⁷

Cell Isolation and Culture

For tumor cell isolation, the proximal joint between the scapula and humerus of wild-type and mutant p22 mice and the adjacent soft tissues were isolated in Dulbecco's phosphate-buffered saline and subjected to a short series of 0.25% trypsin digestions at 37°C to minimize contamination with muscular tissue; after each digestion, the supernatant was discarded. The samples were placed in tissue culture–treated dishes and cut into smaller fragments to promote migration from the segments onto the surface. Cells were grown in Dulbecco's modified Eagle's medium and Gluta-max supplemented with 10% fetal calf serum and 1% penicillin-streptomycin. They were passaged twice before collection to expand the population.

All cells were incubated at 5% CO₂ under humidified atmosphere, and media were changed every 2 to 3 days

before collection. Deletion of *Vhlh* was confirmed by PCR analysis of the genomic DNA extracted from tumor cells and wild-type cells in culture, as previously reported.¹⁵

Western Blotting

Protein extraction and Western blotting for Hif-1 α and Epas1 were performed as previously reported.¹⁶

Results

Loss of Vhl in Mesenchymal Progenitors of the Limb Bud Causes Severe Fibrosis of the Synovial Joints

To investigate the role of Vhl in the biology of mesenchymal progenitor cells, we pursued the conditional deletion of Vhl in limb bud mesenchyme by using Prx1-Cre transgenic mice. We first analyzed *cre* activity in Prx1-Cre mice by taking advantage of the mT/mG reporter.²¹ The mT/mG reporter mice have *loxP* sites on either side of a membrane-targeted tandem dimer tomato (mT) cassette and express red fluorescence in all tissues. When bred to Prx1-Cre mice (PRX1Cre-mTmG), the mT cassette is deleted, allowing expression of the membrane-targeted enhanced green fluorescent protein (EGFP) (mG) cassette in the tissues where *cre* is active. In p14 limbs of

PRX1Cre-mTmG mice, a positive EGFP signal could be clearly detected not only in cartilage, as previously reported,¹⁶ but also in bone, synovium, ligaments, tendons, perimysium, epimysium, and, surprisingly, periendothelial cells within the blood vessel walls, whereas the signal was consistently negative in muscular tissue and endothelial cells (Figure 1).

We next generated *Prx1-Cre;Vhl^{fl/fl}* mutants (VHL) as previously described,¹⁶ and we analyzed the soft tissues that formed and surrounded their synovial joints. VHL mice were born with the expected mendelian frequency and reached adulthood. However, they were smaller than control littermates and died at approximately postnatal day 28 (p28), as already reported.¹⁶

In agreement with our previous findings,¹⁶ analysis of the synovial joints revealed that loss of *Vhl* did not alter either the joint specification (data not shown) or the joint cavitation process in mutant mice when compared with control littermates (Supplemental Figure S1). However, later in fetal development and perinatally, the synovial space in mutant mice appeared to be partially filled with fibrotic tissue (Supplemental Figure S1, B, D, and F). Curiously, the presence of knee menisci was never detected in VHL mutants either prenatally or postnatally (Supplemental Figure S1, C–F, and Figure 2, G and H). After birth, fibrosis of the synovial space was massive in VHL mice, and this was associated with partial destruction of the articular surface cartilage (Figure 2, A–F). The fibrotic lesions were particularly aggressive in the shoulder and in the elbow joints of most mutant mice (Figure 3). These aggressive lesions were formed by numerous spindlyoid cells and a few multinucleated cellular elements; in addition, they had considerable accumulation of an amorphous and acellular material (Figure 3, B and C). Consistent with their fibroblastic appearance, spindlyoid cells within the lesions expressed collagen $\alpha 1$ (Col1a1) mRNA (Figure 3, A–D, G, and H). The aggressive fibrosis eventually penetrated and replaced the growth plate and the adjacent bone (Figure 3, B and H). Fibrotic infiltrations of bone also occurred at the tendon insertions, where they caused erosions and destruction of bony structures, as seen on radiographs (Figure 3, E and F).

Of note, we did not observe fibrosis of the synovial joints in mutant mice in which *Vhl* had been deleted exclusively in cells committed to either the chondrocyte¹³ or the osteoblast lineage^{14,15} (Supplemental Figure S2, A–F) (some data not shown). In agreement with these findings, analysis of *Col2a1-Cre;mTmG^{fl/fl}* (COL2Cre-mTmG) mice revealed only a very low and spotted EGFP signal in the synovium, tendon, and ligaments of the synovial joints (Supplemental Figure S2G).

This finding is consistent with an early mesenchymal population being the target of the *Prx1* enhancer, and it indicates that cells that give origin to the synovial fibrosis are not osteochondroprogenitors.

In addition to the massive fibrosis, loss of *Vhl* in limb bud mesenchyme led to the formation of foci of ectopic cartilage in the soft tissue surrounding the cartilaginous structures




Figure 5 von Hippel Lindau (*Vhl*) deficiency leads to formation of soft tissue tumors in proximity to the synovial joints. **A:** Macroscopic appearance of a postnatal day (p) 28 elbow tumor in VHL mice. The tumor appears as a whitish and multilobulated mass. **B:** Radiograph of a p21 elbow tumor in VHL mice. **C:** Hematoxylin and eosin (H&E) staining of the VHL elbow tumor shown in **B**. The tumor aggressively invades adjacent tissues, such as cartilage, bone, and muscles. **D** and **E:** Higher magnifications of p21 elbow tumor in VHL mice. The tumor is formed by numerous spindlyoid cells with irregular shape and by rare multinucleated cells with an intense eosinophilic cytoplasm indicated by the **yellow arrows** in **E**. **F:** *In situ* cell death detection of a p21 VHL elbow tumor. The **yellow arrows** and the **white arrows** point at terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling–positive cells in the growth plate and in the tumor, respectively. **G:** Radiograph of a p21 shoulder tumor in VHL mice. Note multiple erosions of the mutant olecranon. **H:** H&E staining of the VHL shoulder tumor shown in **G**. An amorphous and acellular material abnormally accumulates in the tumor. Scale bars: 0.5 cm (**A**, **B** and **G**); 100 μ m (**C**–**F** and **H**).

(Figure 2D). These data are in agreement with the notion that hypoxia signaling is not only necessary for chondrogenesis,²⁰ but it may also be sufficient.²⁸

Lastly, fusion of small cartilaginous elements occurred in both forelimb and hindlimb VHL mutant autopods (Figure 2F) (some data not shown). Because no obvious impairment of joint specification could be detected in mutant samples at earlier time points,¹⁶ we concluded that these fusions were not the result of a failure of synovial joints to be specified.

Loss of Vhl in Mesenchymal Progenitors of the Limb Bud Causes Soft Tissue Tumors Located in Close Proximity to Synovial Joints

The aggressive fibrosis described above postnatally was associated with the formation of discrete masses (Figures 4 and 5). The development of these masses mainly occurred in the scapula and elbow (Figure 4, A–J, and Figure 5) and, sporadically, also in the wrist (data not shown), knee (Figure 4, K and L), and sternum (Supplemental Figure S3, B–E). Of note, as with the long bones of the hindlimbs and forelimbs, the sternum originates from the lateral plate mesoderm, as confirmed by *cre* activity at this site²⁹ (Supplemental Figure S3A).

To systematically study these soft tissue lesions, VHL mice and control littermates were analyzed by radiography and histologic analysis at p7 to p28. These masses were overall more frequent in forelimbs than in hindlimbs (Figure 4), which is consistent with the finding that *cre* activity in forelimbs is detectable earlier than in hindlimbs.¹⁹ Radiographs revealed that the VHL masses were partially mineralized and were highly heterogeneous with irregular borders (Figure 4, B–E, G–J, and L, and Figure 5, B and G). Macroscopically, they were whitish and multilobulated (Figure 5A). Histologically, they appeared to aggressively invade adjacent tissues, such as cartilage, bone, and muscles (Figure 5, C and H), and, like the aggressive fibrosis, were formed by numerous spindlyoid cells with irregular shape (Figure 5D) and by rare multinucleated cells with an intense eosinophilic cytoplasm (Figure 5E). Interestingly, the spindlyoid cells did not have any obvious nuclear atypia, and their overall proliferation rate was low (data not shown). Lastly, an amorphic and acellular material abnormally accumulated in the masses (Figure 5, C and H), most likely as a consequence of both cell death (Figure 5F) and synovial fluid being trapped within the lesions.

We next performed an IHC series to further phenotypically characterize the cells that formed these masses. No positive staining for desmin, CD31, von Willebrand factor, CD45, or cytokeratin was detected, which indicates that these cells did not belong to the muscular, endothelial, hematopoietic, or epithelial lineage (Figure 6, A and B, and Supplemental Figure S4, A–C). Conversely, they strongly stained positive for S100 and vimentin (Figure 6, C and D), which are both mesenchymal markers.^{30–34} In addition, they also expressed




Figure 6 von Hippel Lindau (VHL) tumors are mainly formed by Vhl-deficient mesenchymal cells. **A** and **B**: Immunohistochemistry of CD31 (**A**) and desmin (**B**) in VHL tumors. Spindlyoid cells within the tumor are negative for both CD31 and desmin, which indicates that these cells do not belong to either the endothelial or the muscular lineage. **C–F**: IHC of S100 (**C**), vimentin (**D**), and cadherin-11 (**F**) in VHL tumors. **In situ** hybridization for detection of Col1a1 mRNA (**E**). These cells have a mesenchymal phenotype because they strongly stain positive for S100, vimentin, and cadherin-11 and express high levels of Col1a1 mRNA. **G**: Efficient recombination of *Vhl* floxed allele in cells isolated from VHL tumors. Gels have been run under the same experimental conditions. **H**: Western blot analysis of hypoxia-inducible factor (Hif)-1 α (Hif-1 α) and HIF-2 α (Epas1) in protein lysate from CNTRL periarticular soft tissue and from cells isolated from VHL tumors, respectively. Detection of α -tubulin was used as a loading control. Blots were run under the same experimental conditions. Tumor cells have increased accumulation of Hif-1 α and Epas1. **I**: **In situ** hybridization for detection of phosphoglycerate kinase-1 (PGK1) mRNA in a VHL tumor. The tumor expresses high levels of PGK1 mRNA, which is a classic downstream target of Hif-1 α . Scale bar = 100 μ m.




Figure 7 Development of von Hippel Lindau (VHL) tumors is mainly hypoxia-inducible factor (Hif)-2 α (Epas1) independent. Tumors are still detectable in double mutant mice that lack both VHL and Epas1. **A** and **B**: Hematoxylin and eosin (H&E) staining of postnatal day (p) 21 wrists in control (CNTRL) (**A**) and VHL-HIF2 (**B**) mice. **B**: The **black arrow** and the **white arrow** indicate the fibrosis and the fusion of cartilaginous elements, respectively. **C**: Radiograph of a p21 shoulder tumor in VHL-HIF2 mice. **D**: H&E staining of the VHL-HIF2 shoulder tumor shown in **C**. Scale bars: 100 μ m (**A**, **B**, and **D**); 0.5 cm (**C**).

high levels of Col1a1 mRNA (Figure 6E). In light of these data, we thus concluded that the spindlyoid cells that formed the masses had a mesenchymal phenotype.

Notably, numerous spindlyoid cells within the masses stained positively for cadherin-11 (Figure 6F). Cadherin-11 is expressed in fibroblast-like synoviocytes, and it is involved in the development of synovium and tendons.^{35,36} The positive staining results for cadherin-11 suggest that cells within the masses, or at least a subset of these cells, derived from the synovium and/or from tendon progenitors. Interestingly, lack of hemosiderin deposits (data not shown) ruled out the possibility that the masses observed in VHL mice could be the murine equivalent of a human condition known as pigmented villonodular synovitis,³⁷ which is characterized by overgrowth of the synovium, intense inflammation, and accumulation of hemosiderin deposit. In agreement with this conclusion, the virtual absence of CD3-positive staining also excluded the presence of a significant inflammatory component within the lesions (data not shown).

Intriguingly, cells surrounding the necrotic and amorphous areas expressed high levels of secreted phosphoprotein 1 (SPP1) mRNA (Supplemental Figure S4D). SPP1 is involved in numerous pathologic processes, and it is a marker of highly hypoxic tumors.^{38,39}

Taken together, our findings indicate that the masses observed in VHL mutant mice were mainly formed by mesenchymal cells and, despite the lack of nuclear atypia, exhibited features consistent with low-grade malignant

tumors, such as local invasiveness, irregular margins, cell death, presence of multinucleated cells, high cell density, and cellular pleomorphism.

Xenotransplantation has become a widely used tool to determine the tumorigenicity of cells.⁴⁰ However, because it is classically quite challenging to succeed in generating xenografts with tumor fragments or cells isolated from low-grade malignant tumors of the soft tissue,⁴⁰ such an experimental approach was not pursued.

Of note, recombination of the *Vhl* conditional allele in mutant cells isolated from the masses was confirmed by genomic PCR analysis for the recombined (*1-loxP*) allele (Figure 6G). In addition, as predicted, increased accumulation of Hif-1 α and Epas1 proteins occurred in these mutant cells when compared with controls, as indicated by Western blot analysis (Figure 6H). Moreover, most spindlyoid cells within the tumors did not express *Vhl* as indicated by IHC (Supplemental Figure S5). Lastly, consistent with Hif-1 α stabilization, high levels of expression of mRNA encoding for phosphoglycerate kinase-1, which is a classic downstream target of Hif-1 α ,^{20,41} were observed in the VHL masses (Figure 6I).

Soft Tissue Tumors and Synovial Fibrosis Still Develop in Mutant Mice Lacking Both *Vhl* and *Epas1* in Mesenchymal Progenitors of the Limb Bud

In light of our findings, we next asked whether the development of the VHL mesenchymal tumors was *Epas1*




Figure 8 Hypoxia-inducible factor (Hif)-1 α (Hif-1 α) is necessary for the onset of von Hippel Lindau (VHL) tumors. Tumors do not develop in double mutant mice that lack both VHL and Hif-1 α . **A–C:** Radiographs of postnatal day (p) 21 forelimbs in HIF1 (**A**), VHL-HIF1 (**B**), and VHL-HIF1-HIF2 (**C**) mice. **D:** Percentage of VHL, VHL-HIF2, VHL-HIF1, and VHL-HIF1-HIF2 mice presenting tumors detectable by radiography. Scale bar = 0.5 cm (**A–C**).

dependent because it has been proposed that *Epas1* is an oncogene.^{12,42} For this purpose, we generated *Prx1-Cre;Vhl^{fl/fl};Epas1^{fl/fl}* mice, in which both *Vhl* and *Epas1* were conditionally deleted in mesenchymal progenitors of the limb bud (VHL-HIF2 mice). Optimal efficiency of recombination of the *Epas1* conditional allele was confirmed as previously reported.^{16,43} Of note, VHL-HIF2 double mutant mice had an overall postnatal survival similar to VHL mice.

No obvious foci of ectopic cartilage could be detected in double mutants that lack both *Vhl* and *Epas1* (data not shown) at least up to p28, but fusion of small cartilaginous elements still occurred in these double mutant mice (Figure 7, A and B). Synovial fibrosis was ameliorated but not fully corrected by loss of *Epas1* (Figure 7B and Supplemental Figure S6, A, B, F, and G). Notably, loss of *Epas1* rescued menisci formation; however, menisci appeared abnormally fibrotic in VHL-HIF2 mice (Supplemental Figure S6, K and L).

More importantly, VHL-HIF2 mutant mice still developed tumors in proximity to synovial joints (Figure 7, C and D). These masses were phenotypically similar to the ones observed in VHL single mutants as shown by macroscopic inspection, radiographs (Figure 7C), and routine histologic analysis (Figure 7D). However, the overall penetrance of these lesions was 70% in the single VHL mutants but only 47% in the double mutants, as indicated by radiographs at p21 to p28 (Figure 8D).

Notably, whereas tumors in VHL mice had been observed both in proximity to the scapula and at the elbow site with an overall distribution of 55% and 45%, respectively, the totality of the VHL-HIF2 masses was detected exclusively in the scapular region. The biological relevance of this finding is elusive at this stage, but it may be associated with a specific gene expression

profile along the proximal/distal axis of the developing limb.⁴⁴

All in all, these data indicate that *Epas1* is not absolutely necessary for the development of the tumors observed in VHL mutant mice. However, loss of *Epas1* considerably reduced the onset of the tumors in VHL mice. Moreover, *Epas1* is also partially responsible for the fibrosis of their synovial joints as its absence ameliorated the fibrotic infiltration of synovial joints observed in VHL mice.

Development of Soft Tissue Tumors in VHL Mutant Mice Is Hif-1 α -Dependent

Since *Epas1* is not absolutely necessary for the onset of the mesenchymal tumors present in VHL mutant mice, we next studied the involvement of Hif-1 α in this process. For this purpose, we generated *Prx1-Cre;Vhl^{fl/fl};Hif1 α ^{fl/fl}* mice (VHL-HIF1), in which both *Vhl* and Hif-1 α had been deleted in mesenchymal progenitor of the limb bud. These mice were born with the expected mendelian frequency, but, as with VHL mice, premature death occurred at approximately p28, and thus, we were not able to analyze and compare VHL and VHL-HIF1 mutant mice at time points later than p28.

VHL-HIF1 double mutant mice were phenotypically similar to mutant mice lacking only Hif-1 α in limb bud mesenchyme⁴⁵ (HIF1) (Figure 8, A and B). Of note, VHL-HIF1 synovial joints were completely altered and characterized by severe disruption of the articular surfaces and partial fusion of bony epiphysis (Supplemental Figure S6, A, D, F, I, K, and N). Therefore, synovial fibrosis, fusion of small cartilaginous elements, and presence of ectopic cartilage and menisci could not be evaluated in these double mutant mice.

However, we were able to establish that VHL-HIF1 mice did not develop soft tissue tumors at least up to p21 to p28, as shown by X-rays (Figure 8, B and D).




Figure 9 Connective tissue growth factor (Ctgf) is necessary for the onset of von Hippel Lindau (VHL) tumors. Severity of the fibrosis markedly decreases and tumor lesions are not detectable anymore in double mutant mice that lack both VHL and Ctgf. **A:** Immunohistochemistry of Ctgf in a VHL tumor. **B:** Radiographs of a postnatal day (p) 28 VHL-CTGF forelimb. The **black arrow** indicates the initial erosion of the ulna. **C:** Percentage of VHL-CTGF mice presenting tumors detectable by radiography ($n = 18$). Hematoxylin and eosin staining of p28 shoulders (**D** and **E**), p28 elbows (**F** and **G**), and p28 knees (**H** and **I**) in control (CNTRL) (**D**, **F**, and **H**) and VHL-CTGF (**E**, **G**, and **I**) mice. **Black arrows** point to the fibrosis in the synovial space (**E**), infiltration of the fibrosis in the collapsing growth plate (**G**), and menisci (**I**). Scale bars: 75 μm (**A**); 0.5 cm (**B**); 100 μm (**D**–**I**).

To address any potential issue of redundancy and complementarity between Hif-1 α and Epas1, VHL-HIF1-HIF2 triple mutant mice were generated. These mutant mice were born with the expected mendelian frequency, but they also died prematurely around p28. Notably, none of the VHL-HIF1-HIF2 triple mutant mice we studied showed any detectable sign of soft tissue tumors by X-rays (**Figure 8**, C and D). Similarly to HIF1 and VHL-HIF1 mice (**Supplemental Figure S6**, C, D, H, I, M and N), triple mutant mice also

displayed extreme deformities of the limbs and fusion of bony elements; presence of synovial fibrosis and menisci could thus not be completely assessed (**Supplemental Figure S6**, A, E, F, J, K, and O).

Taken together, these findings indicate that Hif-1 α has an essential and nonredundant role in the onset and development of soft tissue tumors in mice that lack Vhl in mesenchymal progenitors of the limb bud.

Development of Soft Tissue Tumors in VHL Mutant Mice Is Ctgf Dependent

It has been reported that both loss of Vhl and stabilization of HIF cause fibrosis through a variety of molecular mechanisms, including up-regulation of profibrotic cytokines such as transforming growth factor- β 1 and Ctgf.⁴⁶

Ctgf (or Ccn2) is a matricellular protein involved in numerous biological processes, including cell proliferation, adhesion, chemotaxis, and extracellular matrix production.⁴⁶ Ctgf is also necessary for proper endochondral bone development.⁴⁷

More importantly, Ctgf is hypoxia inducible in a Hif-1 α -dependent manner,^{46,48} and the expression of Ctgf is up-regulated in a variety of fibrotic disorders.^{49,50}

In light of these findings and in the attempt to understand why Hif-1 α is necessary for the formation of mesenchymal tumors, we analyzed the role of Ctgf in the context of Vhl-deficient limb bud mesenchyme.

IHC confirmed high levels of expression of Ctgf in the VHL tumors (**Figure 9A**). We thus generated *Prx1-Cre;Vhl^{fl/fl};Ctgf^{fl/fl}* (VHL-CTGF) mice lacking both Vhl and Ctgf in limb bud mesenchyme.

VHL-CTGF double mutant mice had an overall postnatal survival similar to VHL single mutants. We studied these double mutants at p28 by radiography and routine histologic analysis. Radiography revealed that none of the VHL-CTGF mice had any evidence of the tumors found in the VHL single mutants (**Figure 9**, B and C). Fibrosis of the synovial joint was still present in both their forelimbs and hindlimbs, but it was considerably milder than in VHL single mutants. In particular, because it occurred in VHL single mutants, histologic analysis revealed that the fibrotic lesions in VHL-CTGF double mutant mice started around tendon/capsule insertions or in the proximity of the synovium (**Figure 9**, D and E), and they invaded the growth plate and the surrounding bony structures, causing erosions detectable by radiography (**Figure 9**, B, F, and G). This fibrotic process acquired a somehow more aggressive phenotype only in a small cohort of VHL-CTGF double mutant mice ($n = 4$ of 18), but it was never associated with the formation of discrete masses.

Of note, as opposed to the VHL mice, the presence of knee menisci was easily identifiable in VHL-CTGF double mutant mice, although their structure was altered by the fibrotic infiltration (**Figure 9**, H and I).

Taken together, these data indicate that loss of Ctgf ameliorated the fibrotic phenotype observed in VHL mutant mice and prevented the formation of mesenchymal tumors at

least up to p28. These findings suggest that the aggressive fibrosis and the formation of discrete soft tissue tumors observed in VHL mutant mice is likely mediated by Hif-1 α through up-regulation of Ctgf.

Discussion

This is the first study reporting the development of severe fibrosis of the synovial joints and Hif-1 α - and Ctgf-dependent soft tissue tumors on loss of Vhl in mesenchymal progenitor cells.

VHL deficiency has been correlated with the development of proliferative lesions in various tissues and organs.^{10,11,51–54} In particular, heterozygous missense mutations of the *VHL* gene have been identified in the VHL syndrome.⁹ This syndrome is characterized by a dominantly inherited tendency to develop pheochromocytoma, clear cell renal carcinoma, and heman-gioblastomas in the central nervous system and retina.⁹ Interestingly, to this end, an augmented predisposition to develop mesenchymal tumors of the soft tissue has not been reported in patients with the VHL syndrome. Our findings, however, suggest that activation of the hypoxia signaling pathway could indeed be an important pathogenetic event in the development of mesenchymal tumors of the soft tissue in humans. To test this possibility, further investigations are warranted.

Notably, loss of Vhl in chondrocytes or in cells of the osteoblast lineage did not cause any fibrosis or tumor of the soft tissue.^{13–15} This finding indicates that the development of fibrosis of the synovial joints and the formation of soft tissue masses in mutant mice that lack Vhl in limb bud mesenchyme are secondary to loss of this protein in an early and unique mesenchymal population, rather than in chondrocytes, osteoblasts, or osteochondroprogenitors (Figure 10). Moreover, our findings of IHC, such as the positive staining results for cadherin-11, suggest that this unique subset of mesenchymal




Figure 10 Up-regulation of Hypoxia-inducible factor (Hif)-1 α (Hif-1 α) and connective tissue growth factor (Ctgf) on loss of von Hippel Lindau (Vhl) in a subset of mesenchymal progenitors leads to fibrosis of the synovial joints and tumors of the soft tissue.

cells could be mainly contributed by the progenitors of tendon, ligament, or synovium cells, although a role for the periendothelial cells cannot be excluded because they also express the *cre* transgene in Prx1-Cre mice.

The tumors characterized in this study are often hypercellular, as indicated by histologic findings. Paradoxically, a significant impairment of proliferation of Vhl-deficient murine embryonic fibroblasts and of growth plate chondrocytes has been previously reported.^{13,55–57} It is thus possible that crosstalk between Vhl-deficient mesenchymal cells and the microenvironment may be critical for the onset and development of the masses observed in VHL mutant mice. Cells in which Hif-1 α and *Epas1* are stabilized through the loss of Vhl could represent a specific niche for additional cell types that may contribute, with yet unknown mechanisms, to the development of the soft tissue masses observed in VHL mutant mice. Along these lines, it is intriguing that multinucleated cells with intense eosinophilic cytoplasm are a consistent histologic feature of the masses we report in this study. The biological relevance of this finding requires further evaluation. In any event, multinucleated cells are often present in malignant mesenchymal tumors of the soft tissue.⁵⁸

A specific role of *Epas1* in the development of renal fibrosis on loss of Vhl has been previously suggested.⁵⁹ In our genetic model, loss of *Epas1* partially ameliorates the massive fibrosis of the synovial space in VHL mutant mice. Our data suggest that *Epas1* overexpression in mesenchymal progenitors, in association with Hif-1 α stabilization, may lead to fibrotic degeneration.

Moreover, our study indicates that *Epas1* is not absolutely necessary for the development of soft tissue tumors on deletion of VHL in limb bud mesenchyme. However, loss of *Epas1* considerably reduced the penetrance of the VHL tumors; thus, *Epas1*, in association with Hif-1 α , may still play a role in the onset of these mesenchymal tumors.

It has been reported that both loss of Vhl and stabilization of HIF cause fibrosis through distinct mechanisms.^{60–64} Along these lines, up-regulation of profibrotic cytokines, such as *Tgfb1* and *Ctgf*,⁴⁶ of inhibitors of metalloproteinases,^{46,65} and of enzymes involved in posttranslational modifications of collagens, such as lysyl oxidase and collagen prolyl-4-hydroxylases,^{46,66,67} have been reported to have an important function in mediating Vhl- or Hif-dependent tissue fibrosis. Notably, *Ctgf* plays profound roles in malignant tumors in major organs and tissues by modulating the actions of key molecules involved in tumorigenesis.⁶⁸ Our study suggests that Hif-1 α -dependent up-regulation of *Ctgf* is the critical event that leads to the formation of fibrosis and soft tissue tumors in the specific developmental setting of synovial joints on loss of Vhl.⁹

Acknowledgments

We thank Drs. M. Celeste Simon, Randall S. Johnson, and Volker H. Haase for providing *Epas1^{fl/fl}*, *Hif1 α ^{fl/fl}*, and *Vhl^{fl/fl}*

mice, respectively; Sirisha Manam for excellent technical support; and the Microscopy and Image-Analysis Lab at University of Michigan for providing outstanding instrumentation.

Supplemental Data

Supplemental material for this article can be found at <http://dx.doi.org/10.1016/j.ajpath.2015.07.008>.

References

- Dunwoodie SL: The role of hypoxia in development of the Mammalian embryo. *Dev Cell* 2009, 17:755–773
- Ivan M, Kondo K, Yang H, Kim W, Valiando J, Ohh M, Salic A, Asara JM, Lane WS, Kaelin WG Jr: HIF1alpha targeted for VHL-mediated destruction by proline hydroxylation: implications for O₂ sensing. *Science* 2001, 292:464–468
- Maxwell PH, Wiesener MS, Chang GW, Clifford SC, Vaux EC, Cockman ME, Wykoff CC, Pugh CW, Maher ER, Ratcliffe PJ: The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. *Nature* 1999, 399:271–275
- Keith B, Johnson RS, Simon MC: HIF1alpha and HIF2alpha: sibling rivalry in hypoxic tumour growth and progression. *Nat Rev Cancer* 2011, 12:9–22
- Haase VH: The VHL tumor suppressor in development and disease: functional studies in mice by conditional gene targeting. *Semin Cell Dev Biol* 2005, 16:564–574
- Kamura T, Sato S, Iwai K, Czyzyk-Krzeska M, Conaway RC, Conaway JW: Activation of HIF1alpha ubiquitination by a reconstituted von Hippel-Lindau (VHL) tumor suppressor complex. *Proc Natl Acad Sci U S A* 2000, 97:10430–10435
- Hsu T: Complex cellular functions of the von Hippel-Lindau tumor suppressor gene: insights from model organisms. *Oncogene* 2012, 31:2247–2257
- Kaelin WG Jr: The von Hippel-Lindau tumour suppressor protein: O₂ sensing and cancer. *Nat Rev Cancer* 2008, 8:865–873
- Kaelin WG: Von Hippel-Lindau disease. *Annu Rev Pathol* 2007, 2:145–173
- Park DM, Zhuang Z, Chen L, Szerlip N, Maric I, Li J, Sohn T, Kim SH, Lubensky IA, Vortmeyer AO, Rodgers GP, Oldfield EH, Lonser RR: von Hippel-Lindau disease-associated hemangioblastomas are derived from embryologic multipotent cells. *PLoS Med* 2007, 4:e60
- Vortmeyer AO, Gnarr JR, Emmert-Buck MR, Katz D, Linehan WM, Oldfield EH, Zhuang Z: von Hippel-Lindau gene deletion detected in the stromal cell component of a cerebellar hemangioblastoma associated with von Hippel-Lindau disease. *Hum Pathol* 1997, 28:540–543
- Kaelin WG Jr: The von Hippel-Lindau tumor suppressor protein and clear cell renal carcinoma. *Clin Cancer Res* 2007, 13:680s–684s
- Pfander D, Kobayashi T, Knight MC, Zelzer E, Chan DA, Olsen BR, Giaccia AJ, Johnson RS, Haase VH, Schipani E: Deletion of Vhlh in chondrocytes reduces cell proliferation and increases matrix deposition during growth plate development. *Development* 2004, 131:2497–2508
- Wang Y, Wan C, Deng L, Liu X, Cao X, Gilbert SR, Boussein ML, Faugere MC, Goldberg RE, Gerstenfeld LC, Haase VH, Johnson RS, Schipani E, Clemens TL: The hypoxia-inducible factor alpha pathway couples angiogenesis to osteogenesis during skeletal development. *J Clin Invest* 2007, 117:1616–1626
- Rankin EB, Wu C, Khatri R, Wilson TL, Andersen R, Araldi E, Rankin AL, Yuan J, Kuo CJ, Schipani E, Giaccia AJ: The HIF signaling pathway in osteoblasts directly modulates erythropoiesis through the production of EPO. *Cell* 2012, 149:63–74
- Mangiavini L, Merceron C, Araldi E, Khatri R, Gerard-O'Riley R, Wilson TL, Rankin EB, Giaccia AJ, Schipani E: Loss of VHL in mesenchymal progenitors of the limb bud alters multiple steps of endochondral bone development. *Dev Biol* 2014, 393:124–136
- Gruber M, Hu CJ, Johnson RS, Brown EJ, Keith B, Simon MC: Acute postnatal ablation of Hif-2alpha results in anemia. *Proc Natl Acad Sci U S A* 2007, 104:2301–2306
- Haase VH, Glickman JN, Socolovsky M, Jaenisch R: Vascular tumors in livers with targeted inactivation of the von Hippel-Lindau tumor suppressor. *Proc Natl Acad Sci U S A* 2001, 98:1583–1588
- Logan M, Martin JF, Nagy A, Lobe C, Olson EN, Tabin CJ: Expression of Cre Recombinase in the developing mouse limb bud driven by a Prxl enhancer. *Genesis* 2002, 33:77–80
- Schipani E, Ryan HE, Didrickson S, Kobayashi T, Knight M, Johnson RS: Hypoxia in cartilage: HIF-1alpha is essential for chondrocyte growth arrest and survival. *Genes Dev* 2001, 15:2865–2876
- Muzumdar MD, Tasic B, Miyamichi K, Li L, Luo L: A global double-fluorescent Cre reporter mouse. *Genesis* 2007, 45:593–605
- Ovchinnikov DA, Deng JM, Ogunrinu G, Behringer RR: Col2a1-directed expression of Cre recombinase in differentiating chondrocytes in transgenic mice. *Genesis* 2000, 26:145–146
- Nagashima T, Kim J, Li Q, Lydon JP, DeMayo FJ, Lyons KM, Matzuk MM: Connective tissue growth factor is required for normal follicle development and ovulation. *Mol Endocrinol* 2011, 25:1740–1759
- Kawamoto T: Use of a new adhesive film for the preparation of multi-purpose fresh-frozen sections from hard tissues, whole-animals, insects and plants. *Arch Histol Cytol* 2003, 66:123–143
- Ohishi M, Ono W, Ono N, Khatri R, Marzia M, Baker EK, Root SH, Wilson TL, Iwamoto Y, Kronenberg HM, Aguila HL, Purton LE, Schipani E: A novel population of cells expressing both hematopoietic and mesenchymal markers is present in the normal adult bone marrow and is augmented in a murine model of marrow fibrosis. *Am J Pathol* 2012, 180:811–818
- Provot S, Schipani E: Fetal growth plate: a developmental model of cellular adaptation to hypoxia. *Ann N Y Acad Sci* 2007, 1117:26–39
- Merceron C, Mangiavini L, Robling A, Wilson TL, Giaccia AJ, Shapiro IM, Schipani E, Risbud MV: Loss of HIF-1alpha in the Notochord Results in Cell Death and Complete Disappearance of the Nucleus Pulposus. *PLoS One* 2014, 9:e110768
- Portron S, Merceron C, Gauthier O, Lesoeur J, Sourice S, Masson M, Fellah BH, Geffroy O, Lallemand E, Weiss P, Guicheux J, Vinatier C: Effects of in vitro low oxygen tension preconditioning of adipose stromal cells on their in vivo chondrogenic potential: application in cartilage tissue repair. *PLoS One* 2013, 8:e62368
- Kaufman MH: The Atlas of Mouse Development. San Diego, CA: Harcourt Brace & Company, 1995, pp. 1–515
- Guarino M, Tosoni A, Nebuloni M: Direct contribution of epithelium to organ fibrosis: epithelial-mesenchymal transition. *Hum Pathol* 2009, 40:1365–1376
- Strutz F, Okada H, Lo CW, Danoff T, Carone RL, Tomaszewski JE, Neilson EG: Identification and characterization of a fibroblast marker: FSP1. *J Cell Biol* 1995, 130:393–405
- Kalluri R, Neilson EG: Epithelial-mesenchymal transition and its implications for fibrosis. *J Clin Invest* 2003, 112:1776–1784
- Franke WW, Grund C, Kuhn C, Jackson BW, Illmensee K: Formation of cytoskeletal elements during mouse embryogenesis. III. Primary mesenchymal cells and the first appearance of vimentin filaments. *Differentiation* 1982, 23:43–59
- Leader M, Collins M, Patel J, Henry K: Vimentin: an evaluation of its role as a tumour marker. *Histopathology* 1987, 11:63–72
- Richardson SH, Starborg T, Lu Y, Humphries SM, Meadows RS, Kadler KE: Tendon development requires regulation of cell condensation and cell shape via cadherin-11-mediated cell-cell junctions. *Mol Cell Biol* 2007, 27:6218–6228
- Lee DM, Kiener HP, Agarwal SK, Noss EH, Watts GF, Chisaka O, Takeichi M, Brenner MB: Cadherin-11 in synovial lining formation and pathology in arthritis. *Science* 2007, 315:1006–1010

37. Mendenhall WM, Mendenhall CM, Reith JD, Scarborough MT, Gibbs CP, Mendenhall NP: Pigmented villonodular synovitis. *Am J Clin Oncol* 2006, 29:548–550
38. Chakraborty G, Jain S, Behera R, Ahmed M, Sharma P, Kumar V, Kundu GC: The multifaceted roles of osteopontin in cell signaling, tumor progression and angiogenesis. *Curr Mol Med* 2006, 6: 819–830
39. Raja R, Kale S, Thorat D, Soundararajan G, Lohite K, Mane A, Karnik S, Kundu GC: Hypoxia-driven osteopontin contributes to breast tumor growth through modulation of HIF1alpha-mediated VEGF-dependent angiogenesis. *Oncogene* 2014, 33:2053–2064
40. Tilokorn D, Daigeler A, Stricker I, Schaffran A, Schmitz I, Steinstraesser L, Hauser J, Ring A, Steinau HU, Al-Benna S: Establishing efficient xenograft models with intrinsic vascularisation for growing primary human low-grade sarcomas. *Anticancer Res* 2011, 31:4061–4066
41. Li H, Ko HP, Whitlock JP: Induction of phosphoglycerate kinase 1 gene expression by hypoxia. Roles of Arnt and HIF1alpha. *J Biol Chem* 1996, 271:21262–21267
42. Shen C, Kaelin WG Jr: The VHL/HIF axis in clear cell renal carcinoma. *Semin Cancer Biol* 2013, 23:18–25
43. Araldi E, Khatri R, Giaccia AJ, Simon MC, Schipani E: Lack of HIF-2alpha in limb bud mesenchyme causes a modest and transient delay of endochondral bone development. *Nat Med* 2011, 17:25–26. author reply 7–9
44. Taher L, Collette NM, Muruges D, Maxwell E, Ovcharenko I, Loots GG: Global gene expression analysis of murine limb development. *PLoS One* 2011, 6:e28358
45. Provot S, Zinyk D, Gunes Y, Kathri R, Le Q, Kronenberg HM, Johnson RS, Longaker MT, Giaccia AJ, Schipani E: Hif-1alpha regulates differentiation of limb bud mesenchyme and joint development. *J Cell Biol* 2007, 177:451–464
46. Higgins DF, Biju MP, Akai Y, Wutz A, Johnson RS, Haase VH: Hypoxic induction of Ctgf is directly mediated by Hif-1. *Am J Physiol Renal Physiol* 2004, 287:F1223–F1232
47. Ivkovic S, Yoon BS, Popoff SN, Safadi FF, Libuda DE, Stephenson RC, Daluiski A, Lyons KM: Connective tissue growth factor coordinates chondrogenesis and angiogenesis during skeletal development. *Development* 2003, 130:2779–2791
48. Nishida T, Kondo S, Maeda A, Kubota S, Lyons KM, Takigawa M: CCN family 2/connective tissue growth factor (CCN2/CTGF) regulates the expression of Vegf through Hif-1alpha expression in a chondrocytic cell line, HCS-2/8, under hypoxic condition. *Bone* 2009, 44:24–31
49. Igarashi A, Nashiro K, Kikuchi K, Sato S, Ihn H, Fujimoto M, Grotendorst GR, Takehara K: Connective tissue growth factor gene expression in tissue sections from localized scleroderma, keloid, and other fibrotic skin disorders. *J Invest Dermatol* 1996, 106:729–733
50. Phanish MK, Winn SK, Dockrell ME: Connective tissue growth factor-(CTGF, CCN2)—a marker, mediator and therapeutic target for renal fibrosis. *Nephron Exp Nephrol* 2010, 114:e83–e92
51. Lee JY, Dong SM, Park WS, Yoo NJ, Kim CS, Jang JJ, Chi JG, Zbar B, Lubensky IA, Linehan WM, Vortmeyer AO, Zhuang Z: Loss of heterozygosity and somatic mutations of the VHL tumor suppressor gene in sporadic cerebellar hemangioblastomas. *Cancer Res* 1998, 58:504–508
52. Lei L, Mason S, Liu D, Huang Y, Marks C, Hickey R, Jovin IS, Pypaert M, Johnson RS, Giordano FJ: Hypoxia-inducible factor-dependent degeneration, failure, and malignant transformation of the heart in the absence of the von Hippel-Lindau protein. *Mol Cell Biol* 2008, 28:3790–3803
53. Couve S, Ladroue C, Laine E, Mahtouk K, Guegan J, Gad S, Le Jeune H, Le Gentil M, Nuel G, Kim WY, Lecomte B, Pages JC, Collin C, Lasne F, Benusiglio PR, Bressac-de Paillerets B, Feunteun J, Lazar V, Gimenez-Roqueplo AP, Mazure NM, Dessen P, Tchertanov L, Mole DR, Kaelin W, Ratcliffe P, Richard S, Gardie B: Genetic Evidence of a Precisely Tuned Dysregulation in the Hypoxia Signaling Pathway during Oncogenesis. *Cancer Res* 2014, 74: 6554–6564
54. Kaelin WG Jr: The von hippel-lindau tumor suppressor protein: an update. *Methods Enzymol* 2007, 435:371–383
55. Mack FA, Rathmell WK, Arsham AM, Gnarra J, Keith B, Simon MC: Loss of pVHL is sufficient to cause HIF dysregulation in primary cells but does not promote tumor growth. *Cancer Cell* 2003, 3:75–88
56. Young AP, Kaelin WG Jr: Senescence triggered by the loss of the VHL tumor suppressor. *Cell Cycle* 2008, 7:1709–1712
57. Young AP, Schlisio S, Minamishima YA, Zhang Q, Li L, Grisanzio C, Signoretti S, Kaelin WG Jr: VHL loss actuates a HIF-independent senescence programme mediated by Rb and p400. *Nat Cell Biol* 2008, 10:361–369
58. Fisher C: *Diagnostic Pathology: Soft Tissue Tumors*. Edited by Osborn EHAG. 1 ed. Salt Lake City, UT: Amirsys Publishing, Inc., 2011, pp. 3. 2-3:106
59. Schietke RE, Hackenbeck T, Tran M, Gunther R, Klanke B, Warnecke CL, Knaup KX, Shukla D, Rosenberger C, Koesters R, Bachmann S, Betz P, Schley G, Schodel J, Willam C, Winkler T, Amann K, Eckardt KU, Maxwell P, Wiesener MS: Renal tubular HIF-2alpha expression requires VHL inactivation and causes fibrosis and cysts. *PLoS One* 2012, 7:e31034
60. Esteban MA, Tran MG, Harten SK, Hill P, Castellanos MC, Chandra A, Raval R, O'Brien TS, Maxwell PH: Regulation of E-cadherin expression by VHL and hypoxia-inducible factor. *Cancer Res* 2006, 66:3567–3575
61. Evans AJ, Russell RC, Roche O, Burry TN, Fish JE, Chow VW, Kim WY, Saravanan A, Maynard MA, Gervais ML, Sufan RI, Roberts AM, Wilson LA, Betten M, Vandewalle C, Bex G, Marsden PA, Irwin MS, Teh BT, Jewett MA, Ohh M: VHL promotes E2 box-dependent E-cadherin transcription by HIF-mediated regulation of SIP1 and snail. *Mol Cell Biol* 2007, 27:157–169
62. Higgins DF, Kimura K, Bernhardt WM, Shrimanker N, Akai Y, Hohenstein B, Saito Y, Johnson RS, Kretzler M, Cohen CD, Eckardt KU, Iwano M, Haase VH: Hypoxia promotes fibrogenesis in vivo via HIF-1 stimulation of epithelial-to-mesenchymal transition. *J Clin Invest* 2007, 117:3810–3820
63. Krishnamachary B, Zagzag D, Nagasawa H, Rainey K, Okuyama H, Baek JH, Semenza GL: Hypoxia-inducible factor-1-dependent repression of E-cadherin in von Hippel-Lindau tumor suppressor-null renal cell carcinoma mediated by TCF3, ZFH1A, and ZFH1B. *Cancer Res* 2006, 66:2725–2731
64. Rankin EB, Tomaszewski JE, Haase VH: Renal cyst development in mice with conditional inactivation of the von Hippel-Lindau tumor suppressor. *Cancer Res* 2006, 66:2576–2583
65. Norman JT, Clark IM, Garcia PL: Hypoxia promotes fibrogenesis in human renal fibroblasts. *Kidney Int* 2000, 58:2351–2366
66. Aro E, Khatri R, Gerard-O'Riley R, Mangiavini L, Myllyharju J, Schipani E: Hypoxia-inducible factor-1 (HIF-1) but not HIF-2 is essential for hypoxic induction of collagen prolyl 4-hydroxylases in primary newborn mouse epiphyseal growth plate chondrocytes. *J Biol Chem* 2012, 287:37134–37144
67. Erler JT, Bennewith KL, Nicolau M, Dornhofer N, Kong C, Le QT, Chi JT, Jeffrey SS, Giaccia AJ: Lysyl oxidase is essential for hypoxia-induced metastasis. *Nature* 2006, 440:1222–1226
68. Kubota S, Takigawa M: Cellular and molecular actions of CCN2/CTGF and its role under physiological and pathological conditions. *Clin Sci* 2015, 128:181–196